arXiv:1207.6357v2 [math.LO] 5 Aug 2012

TWO FAMILIES OF SET THEORIES INTERPRETABLE IN HIGHER ORDER ARITHMETIC

Abstract. A folk theorem says higher order arithmetic has the proof theoretic strength of set theory with limited power set. This paper makes the theorem precise in terms of several axiom systems based on ZF as well as systems based on the elementary theory of the category of sets.

Contents

1. The set theories ZF[n], ZF[n^*], and ZFG[n^*] 1
2. The categorical set theories 3
3. Interpreting ZF[n] in Z_{n+2} 4
References 7

A folk theorem says n-th order arithmetic Z_n has the proof theoretic strength of set theory with a restricted power set, but no precise statement has been published for $n > 2$ and no proof. This paper specifies axioms for ZF[n] based on ZF and for ETR[n] based on the elementary theory of the category of sets (ETCS), each positing that the natural numbers have n successive power sets. In other words each asserts that $\beth_n$ exists where as usual

$$\beth_0 = \aleph_0 \qquad \text{and} \qquad \beth_{n+1} = 2^{\beth_n}$$

Obviously ZF[n] interprets ETR[n] and both interpret Z_{n+2}. We show Z_{n+2} interprets stronger theories ZF[n^*] and ETR[n^*] positing for every set A smaller than $\beth_{n+1}$ and every set B whatever there is a set of all functions $A \to B$. Here to say A is smaller than B, written $A < B$, means there is a one-to-one function $A \to B$. Each of these theories interprets itself plus a global well ordering relation. These extensions are useful in applications as they obviate some appeals to power sets.

1. The set theories ZF[n], ZF[n^*], and ZFG[n^*]

The set theory ZF[0], often called ZF−, is Zermelo-Frankel without power set. The axioms are:

- Extensionality: $\forall z(z \in x \leftrightarrow z \in y) \to x = y$.
- Empty set: $\exists y \forall z \neg[z \in y]$.
- Foundation: $\exists a(a \in x) \to \exists y(y \in x \wedge \neg\exists z(z \in y \wedge z \in x))$.
- Pair set: $\exists w \forall z[z \in w \leftrightarrow (z = x \vee z = y)]$.
- Sum set: $\exists u \forall z[z \in u \leftrightarrow \exists y\, (z \in y \wedge y \in x)]$
- Infinity: $\exists x[\varnothing \in x \wedge \forall y(y \in x \to y \cup \{y\} \in x)]$.
- Replacement: For any formula $\phi(x, y)$ in the language of ZF:

$$\forall x(x \in w \to \exists! y\ \phi(x, y)) \to \exists v\, \forall y\big(y \in v \leftrightarrow \exists x\, (x \in w \wedge \phi(x, y))\big)$$

Examples show how replacement does some things commonly done by power set.

Theorem 1.1. (In ZF[0]) *Any two sets have a cartesian product* $A \times B$*.*

Proof. Form Kuratowski ordered pairs by repeated pair sets. For any sets a, B, replace each $b \in B$ by $\langle a, b\rangle$ to get $\{a\} \times B$. Then replace each $a \in A$ by $\{a\} \times B$ to get $\{\{a\} \times B \mid a \in A\}$ with sum set $A \times B = \{\langle a, b\rangle | \ a \in A, b \in B\}$. □

Theorem 1.2. (In ZF[0]) *Every equivalence relation* $R \subseteq A \times A$ *has a quotient set of equivalence classes.*

Proof. Replace each $a \in A$ by its set of relata $R_a = \{a' \in A | \ \langle a, a'\rangle \in R\}$ to get the set of equivalence classes $A/R = \{\pi \subseteq A | \ \exists a \in A \ (\pi = R_a)\ \}$. □

Theorem 1.3. (In ZF[0]) *There is a unique set* ω *satisfying infinity plus induction, call it the set of natural numbers.*

$$\begin{aligned} &\varnothing \in \omega \wedge \forall y(y \in \omega \to y \cup \{y\} \in \omega) \quad \wedge \\ &\qquad\qquad \forall \sigma \subseteq \omega \ [\forall \ \varnothing \in \sigma \wedge \forall y(y \in \sigma \to y \cup \{y\} \in \sigma) \to \sigma = \omega] \end{aligned}$$

Proof. Given x as in the axiom of infinity take Dedekind's subset:

$$\omega = \{z \in x | \ \forall \sigma \subseteq x \ (\forall \ \varnothing \in \sigma \wedge \forall y(y \in \sigma \to y \cup \{y\} \in \sigma) \to z \in \omega) \ \}$$ □

Theorem 1.4. (In ZF[0]) *Every set* A *has a set* $A^{<\omega}$ *of all finite strings and a set* $\mathrm{Fin}(A)$ *of all finite subsets of* A*.*

Proof. Replacing each $n \in \omega$ by the product set A^n gives $\{A^n | \ n \in \omega\}$ with sum set $A^{<\omega}$. Replacing each n-tuple by the set of its entries gives $\mathrm{Fin}(A)$. □

The ZF[1] axioms further posit ω has a power set $\mathcal{P}(\omega) = \beth_1$, while ZF[2] posits a power set $\mathcal{P}\mathcal{P}(\omega) = \beth_2$, and so on for all ZF[$n$]. Even before seeing the precise axioms of Section 3 it is clear ZF[0] interprets 2nd order arithmetic Z_2, and more generally ZF[n] interprets $(n + 2)$ order arithmetic $Z_{(n+2)}$.

Theorem 1.4 showed ZF[0] implies every set has a set of all its finite subsets. To the contrary, ZF[1] does not imply every set has a set of all its countable subsets. Indeed for no $n > 0$ does ZF[n] imply every set has a set of all countable subsets, let alone all subsets which are smaller than $\beth_n$. Lemma 1.6 shows this.

Lemma 1.5. *ZF[n] for* $0 < n$ *proves these statements are equivalent:*

1. *Every set* A *has a set of all its subsets smaller than* $\beth_n$*.*
2. *For every set* A*, and every* $B < \beth_n$ *there is a set of all functions* $B \to A$

Proof. 1 implies 2 since the graph of a function $B \to A$ is a subset of $B \times A$ the size of B. And 2 implies 1 as every subset of A smaller than $\beth_n$ is the image of some function $B \to A$ with $B < \beth_n$. □

For each $n \in \mathbb{N}$ define ZF[n^*] to be ZF[n] plus either of the conditions in Lemma 1.5. If ZF is consistent then ZF[n] does not imply ZF[n^*] when $n > 0$. Weaker consistency assumptions would suffice but I am not concerned with that:

Lemma 1.6. *ZF proves for every* $1 \leq n \in \mathbb{N}$ *there is a model of ZF[n] where not every set has a set of all its countable subsets.*

Proof. Using ZF and working in L we can assume GCH, so $\beth_n = \aleph_n$. Consider the set $H_{\aleph_\omega + 1}$ of all sets hereditarily smaller than $\aleph_\omega + 1$. This models every ZF[n]. But it also includes $\aleph_\omega$ and all its subsets, while König's inequality shows the set of countable subsets of $\aleph_\omega$ is larger than $\aleph_\omega$. □

On the other hand there is an inner model:

Theorem 1.7. *ZF[n] has a well ordered inner model of ZF[n^*].*

Proof. The inner model L of constructible sets works in ZF[0] nearly as in (Cohen, 1966). Only notice that given L_α (Cohen's M_α) we can get $L_{\alpha+1}$ without using power set, by replacement on the finite strings of elements of L_α and the set of formulas of ZF. Axioms other than power set are verified in Cohen's way and his argument works in ZF[0] to show that if a set A in L has an actual power set then some set in L is a power set for A within L. So ZF[n] proves L is an inner model of ZF[n] and the usual relations are absolute.

Cohen's verification of choice in L defines a well-ordering of L without using power sets, so we can assume choice. As to GCH, Cohen's argument works to show ZF[n] + ($V = L$) proves $\beth_n = \aleph_n$, and that every subset of $\beth_n$ is constructed by some ordinal with cardinality $\beth_n$. So ZF[n] + ($V = L$) proves if $\aleph_{n+1}$ exists it is $\beth_{n+1}$. By cofinality, it proves that when $\aleph_1 < A < \aleph_{n+1}$ then the set of all subsets of A with cardinality less than A is in bijection with A. So ZF[n] shows L restricted to sets hereditarily smaller than $\aleph_{n+1}$ satisfies ZF[n^*]. □

For each $n \in \mathbb{N}$ define ZFG[n^*] to be ZF[n^*] plus a global well-ordering of sets. That is extend ZF[n^*] by a linear order $y \leq_\gamma z$ on sets and an axiom scheme saying every non-empty class has a $\leq_\gamma$-minimal element; and extend the replacement axiom scheme to include formulas using this relation.

Corollary 1.8. *ZF[n] has an inner model of ZFG[n^*].*

2. The categorical set theories

The categorical set theory ETR[0] is ETCS without the axioms of function set or choice, but adding a replacement scheme. Dropping function sets means these axioms do not describe a topos. The axioms are:

- Like a topos: Sets and functions form a category with all finite products, coequalizers, and a subset classifier Ω.
- Nontriviality: Besides $true\colon 1 \to \Omega$ there is a distinct element $false\colon 1 \to \Omega$.
- Infinity: There is a natural number object $\mathbb{N}, 0, s$.
- Extensionality: For any parallel pair of functions $f, g\colon A \to B$ if $f(x) = g(x)$ for all $x \in A$, then $f = g$.
- Replacement: For each relation $R(x, h_1, \ldots, h_n)$ of functions in the language of ETCS this is an axiom:

$$\forall x \in A\ \exists!_i (h_1, \ldots, h_n)\ (R(x, h_1, \ldots, h_n)) \rightarrow \\ \exists (h_1, \ldots, h_n)\ \forall x \in A\ (R(x, x^*h_1, \ldots, x^*h_n)$$

With the axiom of choice, the replacement axiom of McLarty (2004) implies this one. But not without choice. The quantifier $\exists!_i$ reads "exists uniquely up to isomorphism." The idea is that a finite list of functions (meeting some given conditions) gives a finite diagram (of some shape). The notation x^*h for $x \in A$ says h is a function over set A and x^*h its pullback over x:

$$\begin{tikzcd}
x^*S \arrow[rrr] \arrow[drr, "x^*h"] \arrow[ddr] & & & S \arrow[drr, "h"] \arrow[ddr] & & \\
& & x^*T \arrow[rrr] \arrow[dl] & & & T \arrow[dl] \\
& 1 \arrow[rrr, "x"'] & & & A &
\end{tikzcd}$$

The axiom says that if each element of a set A is R-related to a finite diagram (of a given shape) unique up to isomorphism, then there is a diagram (of the same shape) of sets over A such that its pullback over any $x \in A$ is R-related to x.

So separation holds for arbitrary formulas, not only bounded formulas:

Theorem 2.1. (In ETR[0]) *For each property $P(x)$ of arrows this is a theorem:*

$$\forall A\ \exists S\ \exists \text{ a monic } i\colon S \rightarrowtail A\ [\forall x \in A\ (x \in S \leftrightarrow P(x))]$$

Proof. Use replacement on the relation $R(x,h)$ given by

$$(\,P(x)\ \&\ h \cong x \text{ as monics to } A)\ \vee\ (\neg P(x)\ \&\ h \cong \varnothing \to A \text{ as monics to } A) \qquad \square$$

Separation immediately implies every parallel pair of functions $f, g\colon A \to B$ has an equalizer defined by $f(x) = g(x)$. Existence of coequalizers shows every equivalence relation $R \rightarrowtail A \times A$ has a quotient $A \to A/R$. The familiar argument by extensionality in a well-pointed topos works in this context as well to show $true, false$ are the only elements of Ω and their intersection $\varnothing$ has no elements. Separation then shows the empty set $\varnothing$ is initial in the category of sets and functions, Ω is the coproduct $1 + 1$. Replacement on the two elements of $1 + 1$ proves all sets A, B have a coproduct or disjoint union $A + B$.

Separation shows every subset of any set has a Boolean complement, and every two subsets of any set A have an intersection and union. Further, for any relation $r\colon R \rightarrowtail A \times B$ there are subsets of A:

$$\{x \in A | \exists y \in B(R(x,y))\} \quad \text{and} \quad \{x \in A | \forall y \in B(R(x,y))\}$$

Theorem 2.2. (In ETR[0]) *Every set A has a set $A^{<\omega}$ of all finite strings, indeed an $\mathbb{N}$-indexed set $A^{<\omega} \to \mathbb{N}$ assigning each string its length.*

Proof. Essentially like Theorem 1.4. $\square$

The ETR[1] axioms posit $\mathbb{N}$ has a power set $\mathcal{P}(\mathbb{N}) = \beth_1$, while ETR[2] posits a power set $\mathcal{P}\mathcal{P}(\mathbb{N}) = \beth_2$, and so on. The stronger ETR[n^*] posits for any A with a monic $A \rightarrowtail \beth_{n+1}$ and any set B there is an exponential B^A, while ETWO[n^*] adds a global well ordering relation. ZFWO[n^*] obviously interprets all of these.

Clearly ETR[n] interprets Z_{n+2}. So the following section concludes the proof that all of these theories interpret one another.

3. Interpreting ZF[n] in Z_{n+2}

Our n-th order arithmetic Z_n uses a language without product types, and with terms of order $< n$. So first order arithmetic PA$= Z_1$ has natural number terms of order 0. Second order Z_2 adds terms of order 1 for classes of natural numbers, while Z_n has class terms of all orders below n. We write $\forall_i$ or $\exists_i$ to quantify over order

i variables. We adopt the Peano axioms in order 0, except for induction which we state with an order 1 class variable:

$$\forall_1 \sigma\ [(0 \in \sigma \wedge \forall_0 y (y \in \sigma \rightarrow S(y) \in \sigma) \rightarrow \forall_0 y\ (y \in \sigma)\,]$$

We adopt the impredicative comprehension scheme for each order $i < n-1$:

$$\exists_{i+1} X\ \forall x_i\ (x \in X \leftrightarrow \phi(x)\,)$$

for any formula $\phi(x)$ with X not free. This is the system of Simpson (2010, p. 4) extended to higher order, or Takeuti (1987, pp. 192) omitting product types.

3.1. ***j*-indexing.** We adopt any encoding of finite sequences of natural numbers by natural numbers. To encode each (finite or infinite) sequence of classes of order $i > 0$ as a single class of order i we use the idea of j-indexing. Here i is not a term of Z_{n+2} but of the language in which we discuss Z_{n+2}. We cannot quantify on it in Z_{n+2}. In contrast j is an order 0 variable which we can quantify in Z_{n+2}.

Precisely, in Z_{n+2}, use any standard coding of pairs of natural numbers to define indexing operators $\star_i$ by recursion (recursion in our language, not in Z_{n+2}):

$$j \star_0 k \text{ is defined as } \langle j, k\rangle \text{ and } j \star_i x \text{ is defined as } \{j \star_{i-1} y \mid y \in x\}$$

In effect $j \star_i x$ is the class built up from j-indexed numbers just the way x is built up from (un-indexed) natural numbers. Clearly Z_{n+2} proves

$$j \star_i x = j \star_i y \ \rightarrow x = y$$

In Z_{n+2} a *sequence of order i classes* for $i < n+2$, is any order i class where every element is j-indexed for some j. Formally:

$$Seq_i(x) \text{ is defined as } \forall_{i-1} y\ (y \in x \rightarrow \exists_0 j\ \exists_{i-1} y'\ (y = j \star_i y')))$$

The j-component $Comp_i(j, x)$ of any order-i class x is the subclass of x made of j-indexed numbers. Formally:

$$Comp_i(j, x) \text{ is defined as } \{y \in x \mid \exists_{i-1} y'\ (y = j \star_i y')))$$

So a sequence of order i classes is determined by its j-components for all j.

Lemma 3.1. *In any Z_{n+2} and for any $i < n+2$, every sequence of order i classes is the union of its components.*

3.2. **Singleton inclusion.** In Z_{n+2} every order i from 0 to n has a natural image in the next higher. We point out this image is defined by a formula $Sing_i(x, y)$ with variables x, y of orders i and $i+1$ respectively:

$$Sing_i(x, y) \text{ is defined as } \forall_i w\ (w \in y \leftrightarrow w = x)$$

Again i is not a term of Z_{n+2} and we cannot quantifier on it in Z_{n+2}.

Lemma 3.2. *The singleton embedding commutes with j-indexing:*

$$\forall_0 j\ \left(\ Sing_i(x, y) \rightarrow Sing_i(j \star_i x, j \star_{i+1} y)\ \right)$$

So singleton image preserves sequences and their j-components for each j. This will mean that the singleton image of an interpretation of ZF[n] in Z_{n+2} is an interpretation of ZF[n] in Z_{n+3} which we will extend to interpret ZF[$n+1$].

3.3. **The interpretations.** We follow Simpson (2010, §VII.3, pp. 258ff.) interpreting ZF[0] in Z_2. Some exposition will motivate the higher order extension.

3.3.1. *The Z_2 case.* Working in Z_2, interpret a ZF set as a tree, where each node represents a set with one element for each daughter node. In turn a tree is coded as a class of finite sequences of numbers which includes every initial segment of any sequence in it. Each sequence is considered a daughter of its initial segment missing just the last entry. For an example using strings of one-digit numbers take this class of sequences where [] is the empty sequence:

$$T = \{\ [],\ 0,\ 1,\ 10,\ 2,\ 20,\ 21,\ 210\ \}$$

This encodes a downward growing tree:



Three nodes below [] show this represents a three element set. No nodes below the leftmost node 0 shows this node represents the empty set. Altogether this tree encodes the von Neumann ordinal 2, that is $\{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}$. Define the *standard tree* for a finite von Neumann ordinal n as the class of all strictly decreasing sequences of natural numbers that begin with some $j \leq n$.

For the class of all strictly decreasing sequences of natural numbers, each branch remains finite but their lengths are unbounded and the tree is infinitely wide. This represents the von Neumann natural number set $\mathbb{N}$. So the tree interpetation in Z_2 satisfies the axiom of infinity. The empty set axiom is verified by the tree with no nodes beyond the empty sequence [].

We must say which trees represent sets, by means expressible in Z_2:

Definition 3.3. *A class α of sequences of natural numbers is a* suitable tree *if:*

1. *α contains every initial segment of any sequence in α,*
2. *the empty sequence [] is in α,*
3. *it is well-founded: no infinite sequence has all its initial segments in α.*

Consider the suitable classes $\{[], 0, 1\}$ and $\{[], 2\}$ encoding these trees:



At a hasty glance the left tree represents a two element set but both its nodes represent the same set, namely the empty set. We want to construe it as representing the same set $\{\emptyset\}$ as the tree on the right. We need an equivalence relation on suitable classes to represent equality of sets, and we must use this relation to define a relation on suitable classes representing set elementhood.

Simpson (2010, p. 263) handles this in Z_2 and shows the fragment ATR_0 proves these interpret much of ZF[0]. Of course ATR_0 is designedly weaker than replacement but the verification in Z_2 is straightforward: For any relation expressible in Z_2 associating a tree to each daughter of [] in a given tree, Z_2 comprehension suffices to produce a tree replacing each daughter of [] by the corresponding tree.

3.3.2. *Lifting to Z_{n+2}.* The above construction works as well in Z_3 to interpret ZF[0] in order 1 (indeed to give a model, a class of order 2, but we will not use that fact). This interpretation will assuredly not extend to ZF[1] as there are no uncountable classes in order 1. So we adapt definition 3.3 to Z_3:

Definition 3.4. *A class α of sequences of classes of natural numbers is a* suitable Z_3 tree *if:*

1. *α contains every initial segment of any sequence in α,*
2. *the empty sequence [] is in α,*
3. *it is well-founded: no infinite sequence has all its initial segments in α.*

The singleton image of our first interpretation consists of suitable Z_3 trees where all the sequences of classes of natural numbers are sequences of singleton classes. So extend it to all suitable Z_3 trees, with the corresponding interpretation of equality and membership of sets.

This satisfies ZF[0] by the same reasoning as before. To show it satisfies ZF[1] we need some suitable Z_3 tree representing the power set of the natural numbers $\mathbb{N}$. The simplest is the class of all initial sequences of any sequence $\alpha\hat{}\{n\}\hat{}\sigma$ where α is a subset of $\mathbb{N}$, n is in α, and σ is the singleton image of a sequence in the standard Z_2 tree for the von Neumann ordinal n.

The extension to ZF[n] in Z_{n+2} for all n is clear. The chief thing is to replace each natural number in the Z_2 interpretation of ZF[0] by the n-fold singleton of the singleton of ... of that number.

References

Cohen, P. (1966). *Set Theory and the Continuum Hypothesis.* W.A. Benjamin.

McLarty, C. (2004). Exploring categorical structuralism. *Philosophia Mathematica*, pages 37–53.

Simpson, S. (2010). *Subsystems of Second Order Arithmetic.* Cambridge University Press.

Takeuti, G. (1987). *Proof Theory.* Elsevier Science Ltd, 2nd edition.